

**Nevada High School
Youth Risk Behavior Survey (YRBS)
Clark County Comparison Report,
2017-2019**



**School of Community
Health Sciences**

Making Health Happen[™]

Acknowledgements

We would like to acknowledge those who assisted with the 2019 Nevada High School Youth Risk Behavior Survey (YRBS) Report including the Nevada Department of Education; school district superintendents; school principals, teachers and staff; and members of Nevada's Substance Abuse Prevention Coalitions.

Recommended Citation

Young, M., Diedrick, M., Lensch, T., Zhang, F., Peek, J., Clements-Nolle, K., Yang, W. State of Nevada, Division of Public and Behavioral Health and the University of Nevada, Reno. *Nevada High School Youth Risk Behavior Survey (YRBS) Clark County Comparison Report, 2017-2019.*

Funding Source

This surveillance project was partially supported by a grant from the Centers for Disease Control and Prevention (CDC-PS18-1807). Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the CDC.

Additional Resources

UNR YRBS Statewide and Special Reports

<https://www.unr.edu/public-health/research/yrbs>

YRBS Adolescent and School Health

<http://www.cdc.gov/HealthyYouth/yrbs/index.htm>

Nevada YRBS Team Members

State of Nevada, Division of Public and Behavioral Health

Julia Peek, MHA

University of Nevada, Reno School of Community Health Sciences Nevada Center for Surveys, Evaluation, and Statistics

Wei Yang, PhD

Professor and YRBS Principal Investigator

Kristen Clements-Nolle, PhD, MPH

Professor and YRBS Co-Principal Investigator

Meg Diedrick, BS

MPH Graduate Student and Research Assistant

Taylor Lensch, MPH

PhD Graduate Student and YRBS Coordinator

Fannie Zhang, PhD, MS

YRBS Biostatistician

Makena Young

YRBS Intern

Contents

INTRODUCTION	1
METHODS	1
DEMOGRAPHICS	1
Demographic characteristics of participating students	2
2017 TO 2019 COMPARISON	3
Weighted prevalence of participant characteristics and health risk behaviors.....	3
PARTICIPANT CHARACTERISTICS.....	3
VIOLENCE-RELATED BEHAVIORS.....	4
ADVERSE CHILDHOOD EXPERIENCES	5
EMOTIONAL HEALTH.....	6
TOBACCO USE	7
ELECTRONIC VAPOR PRODUCT USE	8
ALCOHOL USE.....	9
MARIJUANA USE	10
OTHER DRUG USE	11
SEXUAL BEHAVIORS	12
DIET.....	13
PHYSICAL ACTIVITY	15
WEIGHT.....	16
OTHER	16

INTRODUCTION

Priority health risk behaviors (i.e. preventable behaviors that contribute to the leading causes of morbidity and mortality) are often established during childhood and adolescence and extend into adulthood. Ongoing surveillance of youth risk behaviors is critical for the design, implementation, and evaluation of public health interventions to improve adolescent health. The Youth Risk Behavior Survey (YRBS) is a national surveillance system that was established in 1991 by the Centers for Disease Control and Prevention (CDC) to monitor the prevalence of health risk behaviors among youth. The Nevada High School YRBS is a biennial, anonymous, and voluntary survey of students in 9th through 12th grade in regular public, charter, and alternative schools. The survey asks students to self-report their behaviors in six major areas of health that directly lead to morbidity and mortality, these include:

- 1) Behaviors that contribute to unintentional injuries and violence;
- 2) Sexual behaviors that contribute to human immunodeficiency virus (HIV) infection, other sexually transmitted diseases, and unintended pregnancy;
- 3) Tobacco use;
- 4) Alcohol and other drug use;
- 5) Unhealthy dietary behaviors; and
- 6) Physical inactivity.

The Nevada High School YRBS provides prevalence estimates for priority risk behaviors and can be used to monitor trends over time.

METHODS

The 2017 and 2019 Nevada YRBS sampling plans were designed to ensure that every eligible student in regular public, charter, and alternative high schools had an equal chance of selection. A random sample of students in grades 9–12 was selected. For the 2017 High School YRBS, 2,019 youth from 49 Clark County schools completed the survey. For the 2019 High School YRBS, 1,989 youth from 49 Clark County schools completed the survey. Please see the 2017 and 2019 Nevada Youth Risk Behavior Clark County Special Reports for more information.

The 2017 and 2019 Nevada YRBS data were weighted based on the sex, race/ethnicity, and grade level of students in each region. The weighting process ensured that the estimates accurately represent the entire student population. Non-response or poor sampling procedures can result in a sample that is not a representative subset of the population; therefore, un-weighted results from these samples may not accurately reflect student behaviors and could be misleading. This report only displays weighted results.

Differences in the weighted prevalence of risk behaviors from 2017 to 2019 were assessed with weighted chi-square tests. Consistent logic checks were applied to allow for comparisons across years. Therefore, estimates may differ from statewide and county reports.

DEMOGRAPHICS

Demographic characteristics of participating students, by sex, age, grade, race/ethnicity, and region — Clark County, Youth Risk Behavior Survey, 2017 & 2019

		2017 Student Participants ^a	2019 Student Participants ^a	2017 Weighted %	2019 Weighted %
Clark County Total		2019	1989	100.00%	100.00%
Sex	Female	1006	1099	48.38%	49.06%
	Male	1000	882	51.62%	50.94%
Age Groups	14 years old or younger	312	276	13.68%	12.82%
	15 years old	499	549	24.15%	25.11%
	16 years old	486	571	25.52%	26.61%
	17 years old	502	411	25.79%	23.35%
	18 years old or older	216	178	10.86%	12.11%
Grade	9th grade	575	545	25.43%	25.99%
	10th grade	479	596	25.63%	26.02%
	11th grade	463	502	24.74%	24.74%
	12th grade	475	341	24.06%	23.18%
	Ungraded or other grade	4	2	0.15%	0.07%
Race/Ethnicity	American Indian/Alaska Native^b	12	9	0.45%	0.41%
	Asian^b	158	178	7.78%	7.90%
	Black^b	168	192	13.54%	13.30%
	Native Hawaiian/Pacific Islander^b	37	38	1.58%	1.68%
	White^b	467	481	26.98%	25.34%
	Hispanic/Latino	999	909	44.05%	45.33%
	Other/Multiple^b	144	138	5.62%	6.04%

^a The sample size in the total and subgroups may differ due to missing data

^b Non-Hispanic

2017 to 2019 Comparison

Weighted prevalence of participant characteristics and health risk behaviors — Clark County, Youth Risk Behavior Survey, 2017 to 2019

PARTICIPANT CHARACTERISTICS

	2017 Percentage and 95% Confidence Interval	2019 Percentage and 95% Confidence Interval	CHANGE Direction and <i>p</i> value
Percentage of high school students who had parents or other adults in the home serving on active duty in the military	4.9% (3.8-6.0)	5.1% (3.9-6.2)	—
Percentage of high school students who qualified for or received free or reduced-price lunches at school	46.2% (41.4-51.0)	46.5% (41.1-51.9)	—
Percentage of high school students who made mostly A's or B's in school during the 12 months before the survey	75.6% (72.4-78.8)	71.1% (67.0-75.2)	—
Percentage of high school students who identified as heterosexual (straight), gay or lesbian, bisexual, or not sure:			
Heterosexual (straight)	83.4% (81.5-85.4)	81.1% (78.8-83.3)	
Gay or lesbian	3.0% (2.3-3.8)	4.0% (3.1-5.0)	
Bisexual	9.9% (8.4-11.4)	10.4% (8.8-12.0)	—
Not sure	3.6% (2.8-4.4)	4.5% (3.5-5.5)	

VIOLENCE-RELATED BEHAVIORS

	2017 Percentage and 95% Confidence Interval	2019 Percentage and 95% Confidence Interval	CHANGE Direction and <i>p</i> value
Percentage of high school students who texted or e-mailed while driving a car or other vehicle during the 30 days before the survey (among students who drove a car or other vehicle)	29.6% (24.9-34.3)	29.0% (22.8-35.2)	—
Percentage of high school students who were threatened or injured by someone with a weapon on school property during the 12 months before the survey (ex. a gun, knife, or club)	7.3% (5.7-8.9)	6.7% (5.2-8.2)	—
Percentage of high school students who were in a physical fight on school property during the 12 months before the survey	7.0% (5.5-8.4)	9.1% (7.1-11.0)	—
Percentage of high school students who were bullied on school property during the 12 months before the survey	15.2% (13.4-16.9)	12.1% (10.4-13.7)	↓ <i>p</i> = 0.0113
Percentage of high school students who were electronically bullied during the 12 months before the survey (including bullied through texting, Instagram, Facebook, or other social media)	11.6% (10.0-13.1)	9.6% (8.1-11.0)	—
Percentage of high school students who did not go to school because they felt unsafe at school or on their way to or from school during the 30 days before the survey	7.9% (6.1-9.6)	13.1% (11.0-15.3)	↑ <i>p</i> = 0.0001
Percentage of high school students who experienced physical dating violence during the 12 months before the survey (including being hit, slammed into something, or injured with an object or weapon on purpose by someone they were dating or going out with; among students who dated or went out with someone during the 12 months before the survey)	7.9% (5.9-9.9)	6.8% (5.3-8.3)	—
Percentage of high school students who experienced sexual dating violence during the 12 months before the survey (including kissing, touching, or physically forced to have sexual intercourse when they did not want to by someone they were dating or going out with; among students who dated or went out with someone during the 12 months before the survey)	9.5% (7.8-11.2)	12.5% (10.3-14.6)	↑ <i>p</i> = 0.0338

ADVERSE CHILDHOOD EXPERIENCES

	2017 Percentage and 95% Confidence Interval	2019 Percentage and 95% Confidence Interval	CHANGE Direction and <i>p</i> value
Percentage of high school students who have ever been hit, beaten, kicked, or physically hurt in any way by an adult (not including spanking for bad behavior)	17.4% (15.5-19.4)	18.5% (16.7-20.3)	—
Percentage of high school students who were ever physically forced to have sexual intercourse when they did not want to	6.8% (5.5-8.1)	5.6% (4.3-6.9)	—
Percentage of high school students who have ever been sworn at, insulted by, or put down by an adult in their home	33.1% (30.7-35.5)	34.9% (32.3-37.5)	—
Percentage of high school students who have ever seen or heard adults in their home slap, hit, kick, punch, or beat each other up	17.1% (15.5-18.7)	18.5% (16.6-20.4)	—
Percentage of high school students who ever lived with someone who was depressed, mentally ill, or suicidal	29.0% (26.7-31.2)	32.3% (30.0-34.6)	↑ <i>p</i> = 0.0410
Percentage of high school students who ever lived with someone who was a problem drinker, alcoholic, or abused street or prescription drugs	31.4% (28.9-33.9)	29.6% (27.0-32.2)	—

2017-2019 Nevada High School YRBS Clark County Comparison Report

EMOTIONAL HEALTH

	2017 Percentage and 95% Confidence Interval	2019 Percentage and 95% Confidence Interval	CHANGE Direction and <i>p</i> value
Percentage of high school students who felt sad or hopeless almost every day for two or more weeks in a row during the 12 months before the survey (so that they stopped doing some usual activities)	34.4% (32.0-36.8)	41.2% (38.5-44.0)	↑ <i>p</i> = 0.0001
Percentage of high school students who seriously considered attempting suicide during the 12 months before the survey	15.9% (14.0-17.8)	17.8% (15.7-19.8)	—
Percentage of high school students who made a plan about how they would attempt suicide during the 12 months before the survey	13.8% (11.9-15.6)	15.4% (13.2-17.5)	—
Percentage of high school students who attempted suicide during the 12 months before the survey	8.2% (6.6-9.9)	8.5% (6.8-10.1)	—
Percentage of high school students who attempted suicide that resulted in an injury, poisoning, or overdose that had to be treated by a doctor or nurse during the 12 months before the survey	2.4% (1.6-3.3)	2.6% (1.7-3.5)	—
Percentage of high school students who did something to purposefully hurt themselves without wanting to die, such as cutting or burning themselves on purpose during the 12 months before the survey	18.0% (16.1-19.8)	20.8% (18.7-22.8)	↑ <i>p</i> = 0.0435
Percentage of high school students who Never/Rarely got the kind of help they need when they felt sad, empty, hopeless, angry, or anxious (among those who felt sad, empty, hopeless, angry, or anxious)	55.2% (52.5-57.9)	55.8% (53.0-58.7)	—

2017-2019 Nevada High School YRBS Clark County Comparison Report

TOBACCO USE

	2017 Percentage and 95% Confidence Interval	2019 Percentage and 95% Confidence Interval	CHANGE Direction and <i>p</i> value
Percentage of high school students who ever smoked cigarettes (even one or two puffs)	21.7% (19.3-24.0)	15.9% (14.1-17.6)	↓ <i>p</i> < 0.0001
Percentage of high school students who smoked cigarettes for the first time before age 13 years (even one or two puffs)	8.7% (7.2-10.2)	5.5% (4.4-6.5)	↓ <i>p</i> = 0.0002
Percentage of high school students who smoked cigarettes during the 30 days before the survey	5.6% (4.3-6.8)	2.7% (1.9-3.4)	↓ <i>p</i> < 0.0001
Percentage of high school students who usually obtained their own cigarettes by buying them in a store or gas station during the 30 days before the survey (among students who currently smoked cigarettes who were aged <18 years)	19.0% (9.9-28.0)	9.9% (0.6-19.1)	—
Percentage of high school students who used smokeless tobacco during the 30 days before the survey	3.1% (2.0-4.2)	3.2% (2.1-4.3)	—

ELECTRONIC VAPOR PRODUCT USE

	2017 Percentage and 95% Confidence Interval	2019 Percentage and 95% Confidence Interval	CHANGE Direction and <i>p</i> value
Percentage of high school students who ever used electronic vapor products (including e-cigarettes, vapes, vape pens, e-cigars, e-hookahs, hookah pens, and mods)	40.6% (36.9-44.3)	40.5% (37.0-43.9)	—
Percentage of high school students who used electronic vapor products during the 30 days before the survey (including e-cigarettes, vapes, vape pens, e-cigars, e-hookahs, hookah pens, and mods)	12.9% (10.5-15.2)	19.2% (16.9-21.6)	↑ <i>p</i> = 0.0001
Percentage of high school students who tried electronic vapor products for the first time before age 13 years (including e-cigarettes, vapes, vape pens, e-cigars, e-hookahs, hookah pens, and mods)	5.8% (4.5-7.2)	6.9% (5.5-8.3)	—
Percentage of high school students who usually obtained their own electronic vapor products by buying them in a store or gas station during the 30 days before the survey (including e-cigarettes, vapes, vape pens, e-cigars, e-hookahs, hookah pens, and mods; among students who currently smoked electronic vapor products and who were aged <18 years)	10.7% (6.5-15.0)	5.7% (3.0-8.5)	↓ <i>p</i> = 0.0440

2017-2019 Nevada High School YRBS Clark County Comparison Report

ALCOHOL USE

	2017 Percentage and 95% Confidence Interval	2019 Percentage and 95% Confidence Interval	CHANGE Direction and <i>p</i> value
Percentage of high school students who ever drank alcohol (not including for religious purposes)	60.0% (56.3-63.6)	55.6% (52.5-58.7)	—
Percentage of high school students who drank alcohol for the first time before age 13 years (other than a few sips)	17.3% (15.1-19.5)	16.7% (14.4-19.0)	—
Percentage of high school students who had at least one drink of alcohol during the 30 days before the survey	24.9% (22.0-27.8)	22.7% (20.5-24.9)	—
Percentage of high school students who participated in binge drinking during the 30 days before the survey (had five or more drinks of alcohol in a row for males, four or more for females within a couple of hours)	13.3% (11.3-15.2)	10.0% (8.6-11.5)	↓ <i>p</i> = 0.0056
Percentage of high school students who rode in a car or other vehicle during the 30 days before the survey driven by someone who had been drinking alcohol	17.2% (15.2-19.3)	13.7% (11.9-15.5)	↓ <i>p</i> = 0.0099
Percentage of high school students who drove a car or other vehicle during the 30 days before the survey when they had been drinking alcohol (among students who drove a car or other vehicle)	5.7% (3.9-7.5)	2.1% (1.1-3.2)	↓ <i>p</i> = 0.0004

2017-2019 Nevada High School YRBS Clark County Comparison Report

MARIJUANA USE

	2017 Percentage and 95% Confidence Interval	2019 Percentage and 95% Confidence Interval	CHANGE Direction and <i>p</i> value
Percentage of high school students who ever used marijuana	34.9% (30.7-39.2)	34.2% (30.7-37.7)	—
Percentage of high school students who tried marijuana for the first time before age 13 years	7.8% (6.2-9.4)	6.7% (5.0-8.4)	—
Percentage of high school students who used marijuana during the 30 days before the survey	18.6% (15.5-21.6)	17.6% (14.9-20.3)	—
Percentage of high school students who usually used marijuana by different methods during the past 30 days before the survey (among those who used marijuana in the past 30 days):			
Smoked it in a joint, bong, pipe or blunt	84.0% (79.4-88.7)	58.1% (51.5-64.6)	
<u>Vaporized it</u>	1.8% (0.0-3.5)	16.9% (12.6-21.2)	↑ <i>p</i> < 0.0001
Used it in some other way	14.2% (10.3-18.2)	25.0% (20.2-29.8)	
Percentage of high school students who rode in a car or other vehicle during the 30 days before the survey driven by someone who had been using marijuana	19.6% (17.0-22.2)	17.3% (14.9-19.6)	—
Percentage of high school students who drove a car or other vehicle during the 30 days before the survey when they had been using marijuana (among students who drove a car or other vehicle)	9.7% (7.1-12.3)	5.7% (4.0-7.4)	↓ <i>p</i> = 0.0058

2017-2019 Nevada High School YRBS Clark County Comparison Report

OTHER DRUG USE

	2017 Percentage and 95% Confidence Interval	2019 Percentage and 95% Confidence Interval	CHANGE Direction and <i>p</i> value
Percentage of high school students who ever used cocaine (any form of cocaine, such as powder, crack, or freebase)	4.3% (3.1-5.5)	4.4% (3.2-5.6)	—
Percentage of high school students who ever used heroin	2.1% (1.2-3.1)	2.4% (1.4-3.4)	—
Percentage of high school students who ever used methamphetamines	2.8% (1.9-3.7)	2.5% (1.5-3.4)	—
Percentage of high school students who ever used ecstasy	5.7% (4.2-7.1)	4.2% (2.9-5.5)	—
Percentage of high school students who ever used synthetic marijuana	6.9% (5.4-8.5)	6.6% (5.1-8.1)	—
Percentage of high school students who ever took prescription pain medicine without a doctor's prescription or differently than prescribed	14.5% (12.5-16.5)	19.2% (17.1-21.2)	↑ <i>p</i> = 0.0014
Percentage of high school students who took prescription pain medicine without a doctor's prescription or differently than prescribed during the 30 days before the survey	6.6% (5.1-8.1)	8.8% (7.3-10.3)	↑ <i>p</i> = 0.0442
Percentage of high school students who ever injected any illegal drug (used a needle to inject any illegal drug into their body)	2.1% (1.3-3.0)	2.1% (1.1-3.1)	—
Percentage of high school students who attended school under the influence of alcohol or other illegal drugs during the 12 months before the survey	15.7% (13.3-18.2)	16.2% (13.7-18.7)	—

2017-2019 Nevada High School YRBS Clark County Comparison Report

SEXUAL BEHAVIORS

	2017 Percentage and 95% Confidence Interval	2019 Percentage and 95% Confidence Interval	CHANGE Direction and <i>p</i> value
Percentage of high school students who ever had sexual intercourse	35.6% (31.4-39.8)	29.7% (26.1-33.4)	↓ <i>p</i> = 0.0335
Percentage of high school students who had sexual intercourse for the first time before age 13 years	3.9% (2.8-5.0)	2.1% (1.3-2.9)	↓ <i>p</i> = 0.0049
Percentage of high school students who had sexual intercourse with four or more persons during their life	9.1% (7.2-11.0)	7.0% (5.3-8.7)	—
Percentage of high school students who had sexual intercourse with at least one person during the 3 months before the survey	24.9% (21.4-28.4)	20.7% (17.7-23.8)	—
Percentage of high school students who drank alcohol or used drugs before last sexual intercourse (among students who were sexually active in the past 3 months)	16.4% (12.5-20.3)	16.3% (12.2-20.5)	—
Percentage of high school students who used a condom during last sexual intercourse (among students who were sexually active in the past 3 months)	56.5% (51.1-61.4)	54.9% (49.2-60.5)	—
Percentage of high school students who used birth control pills before last sexual intercourse (to prevent pregnancy; among students who were sexually active in the past 3 months)	13.6% (9.4-17.8)	17.3% (12.7-21.9)	—
Percentage of high school students who used a shot, patch, birth control ring, IUD, or implant before last sexual intercourse (to prevent pregnancy; among students who sexually active in the past 3 months)	4.6% (2.7-6.5)	5.9% (2.9-8.9)	—
Percentage of high school students who used birth control pills, a shot, patch, birth control ring, IUD or implant before last sexual intercourse (to prevent pregnancy; among students who were sexually active in the past 3 months)	18.2% (13.8-22.7)	23.2% (17.7-28.7)	—
Percentage of high school students who used both a condom and birth control pills, a shot, patch, birth control ring, IUD or implant before last sexual intercourse (to prevent pregnancy; among students who were sexually active in the past 3 months)	5.7% (3.4-8.1)	7.8% (4.3-11.3)	—
Percentage of high school students who <u>did not</u> use any method to prevent pregnancy during last sexual intercourse (among students who were sexually active in the past 3 months)	17.4% (13.1-21.7)	16.9% (12.5-21.3)	—
Percentage of high school students who have ever been pregnant or gotten someone pregnant	1.9% (1.2-2.6)	1.3% (0.8-1.8)	—

2017-2019 Nevada High School YRBS Clark County Comparison Report

DIET

	2017 Percentage and 95% Confidence Interval	2019 Percentage and 95% Confidence Interval	CHANGE Direction and <i>p</i> value
Percentage of high school students who <u>did not</u> eat fruit or drink 100% fruit juices during the 7 days before the survey	6.9% (5.4-8.4)	8.1% (6.7-9.5)	—
Percentage of high school students who ate fruit or drank 100% fruit juices one or more times per day during the 7 days before the survey	52.9% (50.4-55.5)	48.1% (45.7-50.5)	↓ <i>p</i> = 0.0044
Percentage of high school students who ate fruit or drank 100% fruit juices two or more times per day during the 7 days before the survey	25.5% (23.5-27.4)	21.5% (19.4-23.5)	↓ <i>p</i> = 0.0042
Percentage of high school students who ate fruit or drank 100% fruit juices three or more times per day during the 7 days before the survey	16.3% (14.6-17.9)	11.4% (9.9-12.9)	↓ <i>p</i> < 0.0001
Percentage of high school students who <u>did not</u> eat vegetables during the 7 days before the survey	12.4% (10.5-14.3)	12.7% (10.7-14.7)	—
Percentage of high school students who ate vegetables one or more times/day during the 7 days before the survey	9.5% (7.9-11.0)	10.6% (9.1-12.1)	—
Percentage of high school students who ate vegetables two or more times/day during the 7 days before the survey	7.2% (5.6-8.7)	6.5% (5.3-7.7)	—
Percentage of high school students who ate vegetables three or more times/day during the 7 days before the survey	6.6% (5.5-7.8)	4.8% (3.8-5.8)	↓ <i>p</i> = 0.0139
Percentage of high school students who <u>did not</u> drink milk during the 7 days before the survey	26.7% (24.6-28.9)	32.2% (29.6-34.9)	↑ <i>p</i> = 0.0007
Percentage of high school students who drank one or more glasses per day of milk during the 7 days before the survey	25.9% (23.5-28.3)	20.7% (18.7-22.6)	↓ <i>p</i> = 0.0004
Percentage of high school students who drank two or more glasses per day of milk during the 7 days before the survey	14.3% (12.4-16.2)	10.2% (8.7-11.7)	↓ <i>p</i> = 0.0004
Percentage of high school students who drank three or more glasses per day of milk during the 7 days before the survey	6.1% (4.9-7.4)	4.5% (3.5-5.5)	↓ <i>p</i> = 0.0375
Percentage of high school students who <u>did not</u> drink a can, bottle, or glass of soda or pop during the 7 days before the survey (not including diet soda or diet pop)	30.7% (28.2-33.3)	34.6% (32.1-37.0)	↑ <i>p</i> = 0.0318
Percentage of high school students who drank a can, bottle, or glass of soda or pop one or more times per day during the 7 days before the survey (not including diet soda or diet pop)	14.3% (12.3-16.3)	11.4% (9.8-13.0)	↓ <i>p</i> = 0.0237

2017-2019 Nevada High School YRBS Clark County Comparison Report

DIET CONTINUED

	2017 Percentage and 95% Confidence Interval	2019 Percentage and 95% Confidence Interval	CHANGE Direction and <i>p</i> value
Percentage of high school students who drank a can, bottle, or glass of soda or pop two or more times per day during the 7 days before the survey (not including diet soda or diet pop)	9.1% (7.3-11.0)	5.9% (4.8-6.9)	↓ <i>p</i> = 0.0010
Percentage of high school students who drank a can, bottle, or glass of soda or pop three or more times per day during the 7 days before the survey (not including diet soda or diet pop)	5.5% (4.2-6.8)	3.0% (2.1-4.0)	↓ <i>p</i> = 0.0016
Percentage of high school students who <u>did not</u> eat breakfast during the 7 days before the survey	15.9% (14.1-17.7)	14.8% (13.0-16.7)	—
Percentage of high school students who ate breakfast on all 7 days during the 7 days before the survey	32.7% (30.1-35.3)	29.7% (27.1-32.3)	—

2017-2019 Nevada High School YRBS Clark County Comparison Report

PHYSICAL ACTIVITY

	2017 Percentage and 95% Confidence Interval	2019 Percentage and 95% Confidence Interval	CHANGE Direction and p value
Percentage of high school students who <u>did not</u> participate in at least 60 minutes of physical activity on any day during the 7 days before the survey (doing any kind of physical activity that increased their heart rate and made them breathe hard some of the time)	15.1% (13.0-17.2)	18.2% (15.9-20.5)	↑ $p = 0.0423$
Percentage of high school students who were physically active at least 60 minutes per day on 5 or more days during the 7 days before the survey (doing any kind of physical activity that increased their heart rate and made them breathe hard some of the time)	42.8% (39.8-45.8)	35.1% (31.7-38.4)	↓ $p = 0.0005$
Percentage of high school students who were physically active at least 60 minutes per day on all 7 days during the 7 days before the survey (doing any kind of physical activity that increased their heart rate and made them breathe hard some of the time)	22.0% (19.7-24.3)	19.5% (16.9-22.1)	—
Percentage of high school students who watched TV, played video or computer games, or used a computer for 3 or more hours per day (on an average school day; for something that was not school work)	57.9% (55.1-60.8)	61.8% (59.1-64.5)	↑ $p = 0.0481$
Percentage of high school students who attended physical education classes on one or more days in an average school week	53.7% (47.3-60.1)	55.7% (49.2-62.2)	—
Percentage of high school students who attended physical education classes daily (5 days in an average week when they were in school)	24.9% (19.0-30.8)	19.0% (13.7-24.2)	—
Percentage of high school students who played on at least one sports team run by their school or community groups during the 12 months before the survey	47.5% (44.5-50.6)	41.5% (37.8-45.2)	↓ $p = 0.0113$
Percentage of high school students who had 8 or more hours of sleep on an average school night	21.8% (19.9-23.7)	19.8% (17.3-22.3)	—

2017-2019 Nevada High School YRBS Clark County Comparison Report

WEIGHT

	2017 Percentage and 95% Confidence Interval	2019 Percentage and 95% Confidence Interval	CHANGE Direction and <i>p</i> value
Percentage of high school students who were obese (\geq 95th percentile for body mass index, based on sex- and age-specific reference data from the 2000 CDC growth chart)	13.7% (11.7-15.6)	13.0% (11.0-15.0)	—
Percentage of high school students who were overweight (\geq 85th percentile but $<$ 95th percentile for body mass index, based on sex- and age-specific reference data from the 2000 CDC growth chart)	15.4% (13.8-17.0)	15.9% (13.9-17.9)	—

OTHER

	2017 Percentage and 95% Confidence Interval	2019 Percentage and 95% Confidence Interval	CHANGE Direction and <i>p</i> value
Percentage of high school students who Most of the time/Always went hungry because there was not enough food in the house during the 30 days before the survey	3.7% (2.8-4.6)	3.7% (2.8-4.6)	—
Percentage of high school students who visited a dentist during the 12 months before the survey (for a check-up, exam, teeth cleaning, or other dental work)	70.7% (68.1-73.4)	70.0% (67.3-72.6)	—

For additional information regarding this report, contact:

Meg Diedrick, MPH Student
University of Nevada, Reno
School of Community Health Sciences
mdiedrick@unr.edu
775-682-6999

Kristen Clements-Nolle, PhD, MPH
University of Nevada, Reno
School of Community Health Sciences
clements@unr.edu
775-682-7097

<https://www.unr.edu/public-health/research/yrbs>



<http://dpbh.nv.gov/>

